

NaturErhvervstyrelsen orienterer (opdateringsdato 24.11.2016)

I tabellen finder du en liste over planteskadegørere, som kan forekomme i frø eller i urenheder i frøpartiet. Listen er ikke udtømmende, og du vil derfor kunne blive mødt med krav om frihed for andre skadegørere. Du kan se hvilke skadegørere, som er fundet i Danmark/Europa – men specielt for drivhuskulturer kan du risikere at blive mødt med krav om frihed for skadegørere, som normalt ikke forekommer i Danmark/Europa.

Solanum lycopersicon (tomat)

| Skadegører | Frøbåren | Findes i Danmark | Findes i Europa |
|--|----------|---------------------|--------------------|
| Bakterie | | | |
| <i>Bacillus polymyxa</i> | x | | x |
| <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> | x | | x |
| <i>Clavibacter michiganensis</i> subsp. <i>sepedonicus</i> | (x) | | x |
| <i>Pseudomonas corrugata</i> | x | x | x |
| <i>Pseudomonas syringae</i> | x | ? | x |
| <i>Pseudomonas viridiflava</i> | x | | x |
| <i>Ralstonia solanacearum</i> | (x) | | x |
| <i>Xanthomonas campestris</i> pv. <i>raphani</i> | x | | x |
| <i>Xanthomonas gardneri</i> | x | | x |
| <i>Xanthomonas vesicatoria</i> | x | | x |
| | | | |
| Svamp | | | |
| <i>Alternaria alternata</i> | x | x | x |
| <i>Alternaria alternata</i> f.sp. <i>lycopersici</i> | x | | x |
| <i>Alternaria tomato</i> | x | x | x |
| <i>Alternaria solani</i> | x | | x |
| <i>Aspergillus niger</i> | x | | x |
| <i>Athelia rolfsii</i> | x | x | x |
| <i>Chalara elegans</i> (<i>Thielaviopsis basicola</i>) | x | x | x |
| <i>Didymella lycopersici</i> | x | x | x |
| <i>Fulvia fulva</i> (<i>Passalora fulva</i> , tomato leaf mould) | x | x | x |
| <i>Fusarium oxysporum</i> spp. | x | x | x |
| <i>Fusarium</i> spp | x | x | x |
| <i>Gibberella avenacea</i> | x | x | x |
| <i>Gibberella fujikuroi</i> | x | | x |

| | | | |
|--|-----|---|---|
| <i>Glomerella cingulata</i> | x | x | x |
| <i>Golovinomyces orontii</i> | x | x | x |
| <i>Phomopsis</i> sp | x | x | x |
| <i>Phytophthora capsici</i> | x | | x |
| <i>Phytophthora infestans</i> | x | x | x |
| <i>Phytophthora nicotianae</i> (var. <i>parasitica</i>) | x | x | x |
| <i>Pleospora herbarum</i> | x | x | x |
| <i>Pythium</i> sp | x | x | x |
| <i>Septoria lycopersici</i> | (x) | x | x |
| <i>Sclerotinia sclerotiorum</i> | x | x | x |
| <i>Thanatephorus cucumeris</i> | x | x | x |
| <i>Verticillium albo-atrum</i> | x | x | x |
| <i>Verticilium dahliae</i> | x | x | x |
| <i>Verticilium tricorpus</i> | x | | x |
| | | | |
| Virus | | | |
| <i>Alfalfa mosaic virus</i> | x | x | x |
| <i>Arabis mosaic virus</i> | x | x | x |
| <i>Citrus exocortis viroid</i> | | | x |
| <i>Cowpea mild mottle virus</i> | x | | |
| <i>Cucumber mosaic virus</i> | x | x | x |
| <i>Pepino mosaic virus</i> | x | x | x |
| <i>Pepper chat fruit viroid</i> | x | | |
| <i>Potato spindle tuber viroid</i> | x | | x |
| <i>Potato virus Y</i> | | x | x |
| <i>Tobacco mosaic virus</i> | x | | x |
| <i>Tobacco streak virus</i> | x | x | x |
| <i>Tobacco mosaic virus, tomato strain</i> | x | x | x |
| <i>Tomato black ring virus</i> | x | | x |
| <i>Tomato bushy stunt virus</i> | x | | x |
| <i>Tomato mosaic virus</i> | x | | x |
| <i>Tomato ring spot virus</i> | (x) | | x |
| <i>Tomato spotted wilt virus</i> | | | x |
| <i>Tomato yellow leaf curl virus</i> | | | x |
| | | | |
| Skadedyr | | | |
| | | | |